# The Prescribing Patterns and Use of Statins in Patients with **Nonalcoholic Fatty Liver Disease**

## Vandana Khungar<sup>1</sup>, Anna S. Lok<sup>2</sup>, L. Michael Weiss<sup>3</sup>, Paul Thuluvath<sup>4</sup>, Kalyan Ram Bhamidimarri<sup>5</sup>, Derek Gazis<sup>6</sup>, Laura Malahias<sup>6</sup>, K. Rajender Reddy<sup>1</sup>

<sup>1</sup>Division of Gastroenterology and Hepatology, University of Pennsylvania, Philadelphia PA, <sup>2</sup>Division of Gastroenterology and Hepatology, University of Michigan, Ann Arbor MI, <sup>3</sup>Gastro Florida, Clearwater FL, <sup>4</sup>The Institute for Digestive Health & Liver Disease at Mercy, Baltimore MD, <sup>5</sup>University of Miami Health System, Miami, Florida, <sup>6</sup>TARGET PharmaSolutions, Inc. Chapel Hill, NC

# INTRODUCTION

- Among NAFLD patients with metabolic syndrome, the most common cause of death is atherosclerotic cardiovascular disease (ASCVD)
- Although statins are considered safe in liver disease and even potentially beneficial in cirrhosis, many patients are not prescribed a statin due to concerns of hepatic enzyme elevations. • We aim to determine whether statins are under prescribed in NAFLD patients with indications for
- lipid-lowering agents and whether this effect is modified in patients with cirrhosis.

## METHODS

## Cohort

- TARGET-NASH is an ongoing longitudinal, observational cohort of > 3,700 patients with NASH managed according to local practice standards at 55 academic and community sites in the United States.
- Participating clinics provided redacted medical records (structured and unstructured data) from consented patients. Patient narratives, laboratory, pathology, and imaging data were extracted and stored in a secured database. Patient reported outcome (PRO) measures were also collected on a annual basis at select sites. Patients contributed blood samples to a biospecimen repository for biomarker validation and translational research.

## Patient Population

This sub-cohort included 3,284 patients  $\geq$  18 years old with a diagnosis of NAFLD enrolled in TARGET-NASH between August 1, 2016 and October 4, 2018.

Patients were stratified into: NAFLD Cirrhosis, NASH, and NAFL.

## **Clinical Case Definitions of NAFLD**

NAFLD Cirrhosis	History of NAFLD with: 1) Liver biopsy with fibrosis stage = 4 OR 2) Liver biopsy with fibrosis stage = 3 and $1 \ge$ clinical signs of cirrhosis OR 3) 2 or more clinical signs of cirrhosis OR 4) FibroScan® elastography result $\ge$ 11 kPa
	<ul> <li>Confirmed by biopsy:</li> <li>Steatohepatitis by Brunt criteria OR NAS total score ≥ 4</li> <li>Clinical diagnosis:</li> <li>ALT &gt; 19 U/L for adult female (22 child), &gt; 30 U/L for adult male (26 child) and;</li> <li>Hepatic steatoses on biopsy or CT/US/MRI and;</li> <li>I of the following: BMI ≥ 30, type 2 diabetes, dyslipidemia</li> </ul>
NAFL	Any participant not meeting criteria for clinical NASH or cirrhosis

## Statistical Analysis

Indication for statin therapy was defined as one or more of the following:

- Clinical ASCVD
- LDL  $\geq$  160 mg/dL
- Diabetes
- 10-year ASCVD risk at enrollment  $\geq$  7.5%

Statin use was defined as any use documented in the electronic medical record during the follow up period:

**Follow Up Period** 

	Enrollment	
<b>3 years retrospective</b>		1

# RESULTS

- 58.6% of NAFLD patients had one or more indications for lipid lowering therapy
- 61.1% of patients with an indication were prescribed a lipid lowering medication • Men (65.6% vs 59.9%), patients  $\geq$  40 (64.0% vs 31.3%), those with any CV disease history (74.4% vs
- 59.3%) and those with a CV event (79.7% vs 59.4%) were more likely to be prescribed a statin. • Similar proportions of patients managed in academic and community centers, 62.4% and 61.5%, of those with indications were prescribed a statin.
- Statins were used in 65% of those prescribed lipid lowering therapy.

Table 1. Descriptive characteristics of the patients with NAFLD						
Patient Characteristics	NAFLD Cirrhosis (N=1279) N(%)	NASH (N=1317) N(%)	NAFL (N=688) N(%)	p-value		
Statin Status						
Statin Use	480 (37.5)	<b>510 (38.7)</b>	196 (28.5)			
No Statin Use	799 (62.5)	807 (61.3)	492 (71.5)			
Not Available	_	_	_	<0.0001		
Age at Study Entry; Mean (SD)	60.8 (10.2)	53.6 (14.0)	<b>56.1 (13.7)</b>	<0.001		
Sex						
Male	523 (40.9)	816 (62.0)	363 (52.7)			
Not Available	5 (0.39)	6 (0.46)	2 (0.29)	0.0003		
Race						
White	1117 (87.3)	941 (71.5)	374 (54.4)			
Black	34 (2.7)	89 (6.8)	50 (7.3)			
Other	89 (7.0)	222 (16.9)	224 (32.6)			
Not Available	39 (3.0)	65 (4.9)	40 (5.8)	<0.0001		
BMI (kg/m2); Mean (SD)	34.7 (7.5)	33.5 (7.4)	30.3 (7.4)	<0.0001		
<b>History of Coronary Artery</b>	Disease					
Yes	122 (9.5)	66 (5.0)	31 (4.5)			
Not Available	_	_	_	<0.0001		
Diabetes Mellitus						
Yes	875 (68.4)	548 (41.6)	169 (24.6)	<0.0001		
Not Available	_	_	_			
<b>On Acetylsalicylic Acid</b>						
Yes	349 (27.3)	334 (25.4)	147 (21.4)			
Not Available	_	_	_	0.0157		
Smoking Status						
Never	99 (7.7)	94 (7.1)	52 (7.6)			
Current	634 (49.6)	771 (58.5)	403 (58.6)			
Former	376 (29.4)	320 (24.3)	134 (19.5)			
Not Available	170 (13.3)	132 (10.0)	99 (14.4)	<0.0001		



Table 2. Median Laboratory values of patients with NAFLD								
Patient Characteristics	NAFLD Cirrhosis (N=1279)	NASH (N=1317)	NAFL (N=688)	p-value				
Total cholesterol	162	182	183	<0.0001				
HDL	43.0	44.0	48.0	<0.0001				
LDL	90.0	103.0	105.0	<0.0001				
Triglycerides	129.0	148.0	125.0	<0.0001				
AST	38.0	30.0	23.0	<0.0001				
ALT	33.0	41.0	26.0	<0.0001				
HBAlc	6.5	6.2	6.0	0.0007				

### Figure 1. Characteristics of Patients with NAFLD and an Indication for Statin Who Were Prescribed a Statin





# CONCLUSIONS

- particularly in those with decompensated cirrhosis.
- Underprescribing is common among women and younger patients.
- with NAFLD to reduce cardiovascular outcomes.

# ACKNOWLEDGEMENTS

TARGET-NASH is a study sponsored by Target PharmaSolutions (TPS). TPS is a real-world clinical data company based in Durham, NC. The authors would like to thank all the investigators, participants and research staff associated with TARGET-NASH. ClinicalTrials.gov Identifier: NCT02815891

# TARGET-NASH





## 00 EASL for the Study of the Live



• Statins and other lipid-lowering agents are underprescribed in adult patients with NAFLD despite clear indications,

• A shift in treatment paradigm is essential to ensure optimal use of statins and other lipid lowering agents in patients

• For practitioners who still have reservations about statin use, alternative lipid lowering agents may be considered as above.

