Primary Biliary Cholangitis (PBC) in the U.S.: Real World Effectiveness of Obeticholic Acid in TARGET-PBC

Cynthia Levy¹, Christopher L. Bowlus², Elizabeth J. Carey³, George DeMuth⁴, Karen Deane⁴, Marlyn J. Mayo⁵, W. Ray Kim⁶, Bruce R. Bacon⁷, David Bernstein⁸, Paul J. Thuluvath⁹, L. Michael Weiss¹⁰, Uchenna Iloeje¹¹, Mary Erickson¹², Marcie Strauss¹¹, Michael W. Fried¹³

¹University of Miami, Miami, FL, United States ⁴TARGET PharmaSolutions, CA, United States ⁴TARGET PharmaSolutions, Chapel Hill, NC, United States ⁴TARGET PharmaSolutions, CA, United States ⁴TARGET PharmaSolutions, Chapel Hill, NC, United ⁸Northwell Health, Manhasset, NY, United States, ¹²Interept Pharmaceuticals, San Diego, CA, United States, ¹³University of North Carolina Chapel Hill, Chapel Hill, NC, United States ¹⁴Intercept Pharmaceuticals, San Diego, CA, United States, ¹³University of North Carolina Chapel Hill, NC, United States, ¹⁴Intercept Pharmaceuticals, San Diego, CA, United States, ¹⁴University of North Carolina Chapel Hill, NC, United States, ¹⁴Intercept Pharmaceuticals, San Diego, CA, United States, ¹⁴University of North Carolina Chapel Hill, NC, United States, ¹⁴Intercept Pharmaceuticals, San Diego, CA, United States, ¹⁴University of North Carolina Chapel Hill, NC, United States, ¹⁴Intercept Pharmaceuticals, San Diego, CA, United States, ¹⁴Intercept Pharmaceuticals, ¹⁴Intercept Pharmaceuticals, ¹⁴Interce

INTRODUCTION

- Primary biliary cholangitis (PBC) is an uncommon autoimmune cholestatic liver disease. • Up to 40% of PBC patients do not achieve a complete biochemical response to ursodeoxycholic acid (UDCA).
- Obeticholic acid (OCA) was approved by the FDA in 2016 for PBC patients not responding or intolerant to UDCA.

METHODS

- TARGET-PBC is an observational study, initiated in 2016, of adult patients with PBC managed at 25 academic and community Hepatology and Gastroenterology practices.
- Treatment of PBC and initiation of OCA is at the discretion of the treating physician. Patient management will follow each site's local standard of care and no specific treatments, clinical assessments, or laboratory tests will be dictated by enrollment in the study.
- Enrollment after May 2017 was enriched to include only those with elevated serum alkaline phosphatase (ALP) despite UDCA therapy or those treated with OCA or fenofibrate (47% of current enrollment).
- Cirrhosis is derived based on any of the following: stage 4 fibrosis on biopsy, stage 3 fibrosis on biopsy and one secondary condition, two secondary conditions, or stiffness >= 17 on FibroScan. - Secondary conditions include portal hypertension, ascites, varices or collaterals, platelets < 140, cirrhosis on other imaging, and splenomegaly.
- Decompensated requires ascites, encephalopathy, variceal bleeding, or Child–Pugh B or C.

PARTICIPANT CHARACTERISTICS

Total enrollment to date for TARGET-PBC is 405 participants with evaluable data from 63 participants that have ever been prescribed OCA. The age, duration of disease and gender are similar to the baseline characteristics from the POISE trial.

Participant Characteristics at Study Entry (n=63)				
Median Age at Study Entry	56 years			
Mean Duration of Disease	7.7 years (1 to 25 yea			
Gender	92.1% Female (n=5			
Race	81.0% Caucasian (n=			
Ethnicity	20.6% Hispanic (n=)			
AMA Positive	84.7% (n=50)			
Mean Duration of OCA Treatment	10.7 months (1 – 37 mo			
Mean UDCA Dose ¹	16.3 mg/day			
Autoimmune Hepatitis Overlap	19.0% (n=12)			
Cirrhosis	55.6% (n=35)			
Mean Child Pugh Score for Cirrhotic	6.0 (5 – 12)			
Decompensated Cirrhotic	48.6% (n=17)			
Mean Child Pugh Score for Decompensated Cirrhotic	8.0 (5 – 12)			

¹One participant discontinued UDCA due to intolerance

STATEMENT & DISCLOSURES

TARGET-PBC is a collaboration among academic & community investigators, the pharmaceutical industry, and PBC patient community advocates. TARGET-PBC is sponsored by TARGET PharmaSolutions, Inc. TARGET thanks the study staff, nurses, health care providers and patients at each study center for their contributions to this work. Listings of Principal Investigators and Industry Partners are available upon request by emailing info@targetpharmasolutions.com. Cynthia Levy, MD Disclosures: Consultant for Intercept, Novartis, GSK, Cara Therapeutics, TARGET PharmaSolutions; Research Grants from Intercept, Novartis, GSK, Gilead, Genfit, CymaBay, Durect, HighTide, Alnylan, Enanta, Genkyotek.

Mike Fried, MD Disclosures: Dr. Fried receives research grants paid to his institution from AbbVie, BMS, Gilead, Merck, and NIH. He serves as an unpaid consultant to AbbVie, BMS, Merck, and TARGET PharmaSolutions. Stock in TARGET PharmaSolutions is held in an independently managed blind trust.

LABORATORY VALUES

ears) :58) า=51)

nonths)

Last available pre-OCA laboratory values were compared to the last available post start of OCA laboratory values. All laboratory values are based upon site specific local laboratory values.

- 77.4% (n=41) of participants experienced a drop in ALP
- 60.4% (n=32) of participants experienced a drop in ALP >15% - 22.6% (n=12) participants experienced a drop in ALP to < 1.5 ULN
- 81.1% (n=43) of participants experienced a drop in ALT
- 78.8% (n=41) of participants experienced a drop in AST

Laboratory Values	Mean Pre-OCA	Mean Post-OCA
Alkaline Phosphatase (ALP)	381 IU/L (All) 412 U/L (LC) 338 IU/L (Non-LC	265 IU/L (All) 290 IU/L (LC) 233 (Non-LC)
Alanine Aminotransferase (ALT)	62.0 IU/L (All) 69.5 IU/L (LC) 51.4 IU/L (Non-LC)	48.0 IU/L (All) 47.3 IU/L (LC) 49.0 IU/L (Non-LC)
Aspartate Aminotransferase (AST)	62.9 IU/L (All) 74.7 IU/L (LC) 46.8 IU/L (Non-LC)	52.1 IU/L (All) 57.6 IU/L (LC) 45.2 (Non-LC)
Total Bilirubin (TB)	1.45 mg/dL (All) 1.90 mg/dL (LC) 0.84 mg/dL (Non-LC)	1.63 mg/dL (All) 2.03 mg/dL (LC) 1.13 mg/dL (Non-LC)
Platelets	204.4 10^3/uL (All) 171.5 10^3/uL (LC) 250.1 10^3/uL (Non-LC)	188.8 10^3/uL (All) 143.1 10^3/uL (LC) 245.4 10^3/uL (Non-LC)
Albumin	4.0 g/dL (All) 3.8 (LC) 4.1 (Non-LC)	3.8 g/dL (All) 3.6 g/dL (LC) 4.1 g/dL (Non–LC)
Creatinine	0.73 mg/dL (All) 0.72 mg/dL (LC) 0.73 mg/dL (Non–LC)	0.73 mg/dL 0.72 mg/dL (LC) 0.74 (Non–LC)
INR	1.02 (All) 1.03 (LC) 0.97 (Non-LC)	1.09 (All) 1.14 (LC) 1.00 (Non-LC)


LC = Liver Cirrhosis

PARTICIPANTS WITH ABNORMAL BILIRUBIN

Most participants (42/52, 80.8%) had normal total bilirubin (TB) prior to OCA start and remained essentially unchanged during treatment. Ten participants (19.2%) had elevated TB prior to start of OCA.

	Pre-OCA TB	Post-OCA TB	Continued Treatment	Current OCA Dose	Listed For Transplant	Reason for OCA Discontinuation
1	2.0 mg/dL	2.0 mg/dL	Yes	5.0 mg QD	No	
2	2.1 mg/dL	1.5 mg/dL	No		No	Discontinued OCA due to pruritus
3	2.4 mg/dL	2.3 mg/dL	No		No	Discontinued OCA due to pruritus
4	2.5 mg/dL	0.7 mg/dL	Yes	5.0 mg QD	No	
5	2.6 mg/dL	4.2 mg/dL	Yes	10 mg BIW	No	
6	2.6 mg/dL	4.8 mg/dL	No		No	Discontinued OCA due to elevation in TB
7	3.5 mg/dL	11.5 mg/dL	No		No	Discontinued OCA due jaundice and potential for acute drug injury
8	3.6 mg/dL	4.9 mg/dL	No		No	Discontinued OCA due to lack of efficacy
9	9.6 mg/dL	7.2 mg/dL	No		No	Discontinued OCA due to pruritus
10	10.6 mg/dL	15.1 mg/dL	Yes	5.0 mg BIW	No	

OBETICHOLIC ACID DOSING



ADVERSE EVENTS*

Adverse events are defined as any new medical event, regardless of causality, captured in the medical records after the start of OCA.

Pruritus 25.4% (n=)

Fatigue 12.7% (n=8)

Abdominal Pain 9. (n=6)

Constipation 6.3%

Osteopenia 6.3% (

Urinary Tract Infect 6.3% (n=4)

CULLICIONC GUNGLUJIUNJ

- of all patients had decompensated cirrhosis. • The results in this more advanced population are consistent with the POISE RCT results and use of OCA had a beneficial effect on the biochemical endpoints.
- TARGET-PBC participants with the most advanced disease (BL bilirubin over 2mg/DL), 5 improved/stayed the same; 5 worsened/stayed the same; none advanced to LT.
- In participants without cirrhosis, platelet count and albumin tended to stay stable, but in participants with cirrhosis trended towards deterioration, suggesting that earlier intervention before cirrhosis might be more beneficial to slow progression.
- The frequency of pruritus was lower than reported in the POISE clinical trial. However, the rate of treatment discontinuation we reported is higher, likely because of the more advanced disease population in this cohort

P-Valu	e
<.0001 (A <.0001 (L .0030 (Non	C)

Mean % Change

-21.5% (All)

-22.7% (LC)

-20.0% (Non-LC)

-18.8% (All)

-26.3% (LC)

-8.16% (Non-LC)

-14.5% (All)

-21.0% (LC)

-5.62% (Non-LC)

4.29% (All)

-4.12% (LC)

15.77% (Non-LC)

-6.3% (All)

-12.5% (LC)

2.2% (Non-LC)

-3.59% (All)

-5.37% (LC)

-1.04% (Non-LC)

10.05% (All)

17.73% (LC)

-1.05% (Non-LC)

6.65% (All)

8.18% (LC)

3.02% (Non-LC)

<.0001 (All) <.0001 (LC) .1433 (Non-LC) <.0001 (All) <.0001 (LC) .1528 (Non-LC)

.9189 (All) .4354 (LC) .3833 (Non-LC)


.0095 (All) .0003 (LC) .9661 (Non-LC)

.0700 (All) .0325 (LC) .8730 (Non-LC)

.7060 (All) .8124 (LC) .7819 (Non-LC)

> .2410 (All) .3470 (LC)

.8750 (Non-LC)





Lack of Efficacy

=16)	Back pain 4.8% (n=3)	Depression 3.2% (n=2)	Oedema peripheral 3.2% (n=2)
3)	Dry Eye 4.8% (n=3)	Elevated liver enzymes 3.2% (n=2)	Pyrexia 3.2% (n=2)
.5%	Insomnia 4.8% (n=3)	Haemorrhoids 3.2% (n=2)	Vertigo 3.2% (n=2)
(n=4)	Gastroesophageal reflux disease 4.8% (n=3)	Hypokalaemia 3.2% (n=2)	Vitamin D deficiency 3.2% (n=2)
n=4)	Anaemia 3.2% (n=2)	Nasopharyngitis 3.2% (n=2)	Vomiting 3.2% (n=2)
ction	Arthralgia 3.2% (n=2)	Nausea 3.2% (n=2)	

1.6% (n=1)

 The participants in the TARGET-PBC cohort were more advanced than those studied in POISE, the randomized clinical trial (RCT) for OCA approval; here, 19% had AIH overlap, >50% had Cirrhosis, almost a third

 Better education of providers on management of pruritus while on OCA may improve drug tolerance and help reduce drug discontinuation.